

# **California Bioscience**

83103 Avenue 48, Ste.1B #204 Coachella, CA 92236 USA Phone : +1.6268339877 Email : info@cali-bio.com

# **Product Datasheet**

Product Name	Cystatin C Rat Recombinant
Cata No	CB501058
Source	Escherichia Coli.
Synonyms	Cystatin-C, Cystatin-3, CYSC, MGC105556.

### Description

Cystatins are a superfamily of cysteine proteinase inhibitors found in both plants and animals. They comprise a group of proteinase inhibitors, widely distributed in tissues and body fluids, and form tight complexes with cysteine proteases such as cathepsin B, H, L and S. Cystatin C, a secreted molecule of this family, is of interest from biochemical, medicine and evolutionary points of view. Cystatin C, with molecular weight of 13260 Da, is composed of 120 amino acids, lacks carbohydrate and has two disulfide bridges located near the carboxyl terminus. Cystatin C is increased in patients with malignant diseases, and is related to the insufficiency of renal function and appears to be a better marker than creatinine. On the other hand, low levels of cystatin C involve cause the breakdown of the elastic laminae and, subsequently, the atherosclerosis and abdominal aortic aneurysm. Total 134 AA, Mw: 14.93 kDa (calculated). N-terminal His-tag (14AA).

## **Physical Appearance**

Sterile Filtered White lyophilized (freeze-dried) powder.

#### **Purity**

Greater than 96% as determined by SDS-PAGE.

# Formulation

Filtered (0.4 micron) and lyophilized from 0.5 mg/ml in 0.03M Acetate buffer, pH 4.

#### Reconstitution

Add 0.2 ml of 0.1M Acetate buffer pH4 and let the lyophilized pellet dissolve completely. For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 0.1mg/ml. In higher concentrations the solubility of this antigen is limited.

#### Stability

Store lyophilized protein at -20 $^{\circ}$ C. Aliquot the product after reconstitution to **avoid repeated freezing/ thawing cycles**. Reconstituted protein can be stored at 4 $^{\circ}$ C for a limited period of time; it does not show any change after two weeks at 4 $^{\circ}$ C. Minimum shelf life 2 years.

#### Sequence

MRGSHHHHHH GMASGTSRPP PRLLGAPQEA DASEEGVQRA LDFAVSEYNK GSNDAYHSRA IQVVRARKQLVAGINYYLDV EMGRTTCTKS QTNLTNCPFH DQPHLMRKAL CSFQIYSVPW KGTHTLTKSS

## Applications

Western blotting.

# \* For Non-Clinical Research Use Only \*